

## Durham Research Online

---

### Deposited in DRO:

27 February 2009

### Version of attached file:

Accepted Version

### Peer-review status of attached file:

Peer-reviewed

### Citation for published item:

Macpherson, C. G and Dreher, S. T and Thirlwall, M. F (2006) 'Adakites without slab melting : high pressure differentiation of island arc magma, Mindanao, the Philippines.', *Earth and planetary science letters.*, 243 (3-4). pp. 581-593.

### Further information on publisher's website:

<http://dx.doi.org/10.1016/j.epsl.2005.12.034>

### Publisher's copyright statement:

### Additional information:

---

### Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

**Adakites without slab melting: high pressure differentiation of  
island arc magma, Mindanao, the Philippines**

Colin G. Macpherson<sup>1,\*</sup>, Scott T. Dreher<sup>1</sup>, Matthew F. Thirlwall<sup>2</sup>

<sup>1</sup> Department of Earth Sciences, University of Durham, Durham, DH1 3LE, UK.

<sup>2</sup> Department of Geology, Royal Holloway University of London, Egham, TW20 OEX  
UK.

\* Corresponding Author

E-mail: colin.macpherson@durham.ac.uk

Tel: +44 (0)191 334 2283

Fax: +44 (0)191 334 2301

Accepted for publication by *Earth and Planetary Science Letters* 23 December 2005

Word Count	Abstract	208
	Text	4552
	References	2099 (72 citations)
	Figure Captions	669
	+ 6 Figures and 2 Tables of Supplementary Data.	

## 22 ABSTRACT

23 New geochemical data for Pleistocene magmatic rocks from the Surigao peninsula,  
24 eastern Mindanao, the Philippines, demonstrate typical adakitic traits, including elevation  
25 of Sr/Y and depletion of the heavy rare earth elements.  $^{87}\text{Sr}/^{86}\text{Sr}$  and  $^{143}\text{Nd}/^{144}\text{Nd}$  ratios of  
26 the adakites do not support melting of the subducted Philippine Sea Plate but resemble  
27 Pliocene arc lavas generated in the same subduction zone. Excepting the heavy rare earth  
28 elements, the adakites and arc lavas also possess similar ratios of incompatible elements  
29 suggesting that the adakites were ultimately derived from melting of the mantle wedge.  
30 The wide range of  $\text{SiO}_2$  in the adakites and its strong correlation with trace element  
31 concentrations and ratios indicate two possible mechanisms for generating the adakitic  
32 signature. (1) Adakitic melt was produced from basaltic arc magma by fractional  
33 crystallisation of a garnet-bearing assemblage. (2) Solidified basaltic rock containing  
34 garnet melted to yield adakitic magma. In either case the basaltic precursor was generated  
35 from fluid-modified mantle then differentiated within the garnet stability field. In Surigao  
36 this requires differentiation within mantle. The Surigao example suggests that any  
37 subduction zone has the potential to produce adakitic magma if basalt crystallises at  
38 sufficient depth. This has important implications for the geodynamics of modern and  
39 ancient subduction zones that have generated similar rocks.

40 Keywords: subduction; magmatism; differentiation; adakite; slab

## 41 1. Introduction

42 Thermal models predict that hydrated basalt in subducted ocean crust is too cold to melt  
43 when it lies beneath the volcanic arc of most modern subduction zones [1,2]. While some  
44 models incorporate melting of subducted crust [3], the geochemistry of arc lavas indicates  
45 (i) that devolatilisation is the main mechanism transferring material out of the slab, and  
46 (ii) that the overlying mantle wedge is, volumetrically, the major source of arc lavas [4-  
47 7]. Partial melting of subducted crust should leave a garnet-bearing residue [8,9],

48 producing magmas with intermediate  $\text{SiO}_2$ , elevated  $\text{Al}_2\text{O}_3$ ,  $\text{Sr/Y}$  and  $\text{La/Y}$ , and low  $\text{Y}$ .  
49 Rocks of this type, which have become known as adakites, have been generated in active  
50 subduction zones where young ocean crust is subducted ( $<25\text{Ma}$ ). This observation has  
51 been interpreted as evidence that young ocean crust is more prone to melting than older  
52 crust because it retains a greater proportion of its initial heat [10].

53 Constraining the origin of modern adakites is important for several reasons. First, the  
54 presence of adakitic rocks implies an unusual thermal regime compared to most modern  
55 subduction zones. Second, many major and trace element characteristics of adakites  
56 resemble tonalite-trondhjemite-granodiorite gneisses, which are important components in  
57 Archean terranes. Therefore, modern adakitic magmatism may provide an analogue for  
58 continental growth processes in the early Earth [11-14]. Finally, several suites of adakitic  
59 rocks are associated with porphyry and epithermal style  $\text{Cu}$ -,  $\text{Au}$ -mineralization [15-17].  
60 While the metallogenic significance of this link is contested [18-20] the association offers  
61 to shed light on the thermal and dynamic state of subduction zones that host such  
62 deposits.

63 Since their first description as products of melting young slab [10], increasing numbers of  
64 adakitic suites have been recognized that were emplaced where the subducted crust was  
65 old, and thus inferred to be cold. This observation has two possible implications for the  
66 slab-melting hypothesis. The first possibility is that certain exceptional subduction zone  
67 geometries permit melting of subducted basaltic rocks which are greater than 25 million  
68 years of age. Several such mechanisms have been advanced including melting the leading  
69 edges of newly subducted slabs [21], shear heating of slab interiors that are exposed along  
70 fracture zones [22], or prolonged slab residence in the shallow mantle as a result of  
71 decreasing angle of subduction [23]. Each of these “cool slab” models appeals to a unique  
72 thermal structure and slab melting mechanism for the subduction zone in question.

73 The alternative implication is that melting of subducted crust does not generate all, or  
74 even any, adakitic magma. For example, arc crust that is sufficiently thick for garnet to be  
75 stable in basaltic rock is proposed as a source for adakitic magmatism in the Andes,  
76 western US and Tibet [24-28]. However, this mechanism is not feasible where arc crust is  
77 less than ~30km thick; the minimum depth of garnet amphibolite or eclogite P-T  
78 conditions. For arcs with thin crust this has led to the default interpretation that subducted  
79 crust is the only part of the subduction zone where basaltic rocks can attain a suitable  
80 mineralogy to act as adakite sources.

81 The Surigao peninsula in Mindanao, the Philippines, hosts adakitic rocks generated  
82 during subduction of the Philippine Sea Plate at the Philippine Trench (Fig. 1). This plate  
83 margin initiated at ~ 7Ma, to the east of Luzon, since when it has propagated southwards.  
84 Subduction of the Philippine Sea Plate beneath Mindanao began in the late Miocene or  
85 early Pliocene [29]. The Philippine Sea Plate crust that was subducted beneath Mindanao  
86 at that time was more than 50 million years old [30] and so was too cold to melt under  
87 normal subduction zone conditions [1]. Furthermore, the Surigao crust is relatively thin  
88 and unlikely to host garnet-bearing basaltic rocks [21]. To account for the presence of  
89 adakites by slab melting, Sajona et al. [21] required a mechanism to melt the cool slab. In  
90 any incipient subduction zone a large thermal contrast will exist between the leading edge  
91 of the new slab and the mantle it penetrates. Numerical simulations indicate that the  
92 leading edge of a new slab may be heated to melting point, even if it is more than 25  
93 million years old [1]. Therefore, Sajona et al. [21] proposed that the presence of  
94 Pleistocene adakites in Surigao indicates an old slab melting in a very young subduction  
95 zone, rather than melting of a young slab.

96 We present new geochemical data to test the slab-melt hypothesis in Surigao. Our data  
97 indicate that the geochemical distinction between these adakites and more normal island  
98 arc lavas (generated by the same subduction zone) result from differentiation of the  
99 adakites in the garnet stability field. Our model removes the need to postulate several

different mechanisms to melt old subducted crust. Instead, adakites can be regarded as part of the spectrum of magmas that may be produced by any subduction zone.

## 2. Surigao del Norte

Surigao del Norte lies at the northern extremity of eastern Mindanao (Fig. 1a). The basement consists of ophiolitic rocks overlain by volcanic and sedimentary rocks derived from a Palaeogene arc [31,32]. In the east, the mountains of the Pacific Cordillera rise towards the south while the western peninsula is dominated by the elongate Malimono Ridge (Fig. 1b and c). Between these is a low-lying central plain occupied in the south by Lake Mainit. The plain is separated from the Malimono Ridge by the Philippine Fault, a sinistral strike-slip fault extending the length of the Philippine archipelago (Fig. 1a and b). To the east, the central plain meets the Pacific Cordillera along Oligo-Miocene reverse faults that show evidence of recent reactivation as normal faults (Fig. 1b). Since initiation of the Philippine Fault during the early Pleistocene the central plain has evolved as a down-faulted basin [32]. Extension, and implied lithospheric thinning, to produce the basin could result from transtension across an eastward step in the Philippine Fault south of Surigao del Norte [32], from trench-normal extension due to rollback of the new slab [33], or from trench-parallel extension due to oblique subduction of the Philippine Sea Plate [34].

\_\_\_\_\_ (Figure 1 to be close to this section) \_\_\_\_\_

Magmatic rocks associated with the Philippine Trench are preserved as hyababyssal stocks and lava flows throughout the central and western part of Surigao del Norte (Fig. 1b). Intrusive relationships with sedimentary units and radiometric dating [31] indicate that magmatism in the eastern part of the peninsula occurred from the very latest Miocene into the Pliocene. Magmatism from the Quaternary cone of Mt. Maniayao and further to the west (Fig. 1b) is Pleistocene or younger [31]. Major and trace element concentrations and

Sr and Nd isotopic ratios were determined for a new suite of rocks that includes both Pliocene and Pleistocene suites (Fig. 1b; see Supplementary Data).

### 3. Results

Late Miocene to Pliocene rocks, hereafter called the Pliocene arc, are island arc basaltic andesite and andesite (Fig. 2, 3 and 4a). The Pleistocene igneous rocks display characteristics typical of adakites, such as intermediate  $\text{SiO}_2$  contents, elevated  $\text{Al}_2\text{O}_3$ , high Sr/Y, and low Y and heavy rare earth element (HREE) concentrations (Table 1; Figs. 2-4). However, Sr is not the only incompatible trace element that is enriched relative to Pliocene rocks with similar  $\text{SiO}_2$ ; all elements except Y and the HREEs are enriched by similar amounts (Fig. 4a,b). Within the Pleistocene suite it is the depletion in Y that is the major control on the development of the adakitic signature (Fig. 2d,e, Fig. 4a). The strength of the adakite signature is highly variable within the suite and correlates with silica (Figs. 3a-c and 4b). There is also a geographic control on composition; the adakitic signal is strongest in the west and decreases towards Mt. Maniayao (Fig. 3e and f).

\_\_\_\_\_ (Figures 2, 3 and 4 to be close to this section) \_\_\_\_\_

$^{87}\text{Sr}/^{86}\text{Sr}$  and  $^{143}\text{Nd}/^{144}\text{Nd}$  display similar, limited ranges in the Pleistocene and Pliocene rocks. Both are distinct from the upper mantle beneath Mindanao, which is believed to resemble I-MORB [35], and from the composition of unleached, altered basalt drilled from the Philippine Sea Plate immediately outboard of the Philippine Trench (Fig. 5).

### 4. Discussion

The strength of the adakite signature in Pleistocene rocks shows significant variation. Large ranges in Sr/Y are common to several adakitic suites [e.g. 10,21,24,36], however, the correlation with, and wide range of,  $\text{SiO}_2$  suggests that the adakite signal at Surigao (i) was diluted by a more mafic component, (ii) varied in response to changing degrees of slab melting, or (iii) developed from a more mafic, arc-like magma or basaltic protolith. If

the last possibility is true then adakitic magmas can be developed without the need to invoke slab melting.

#### *4.1 Modification of Adakitic Slab Melts*

Two mechanisms could modify the composition of true slab melts towards those of arc lavas. Slab melts could mix with contemporaneous arc lavas or they could interact with mafic or ultramafic rocks during transport from their source to the surface.

Mixing between a strongly adakitic magma and a more mafic island arc magma (low SiO<sub>2</sub>, Sr/Y and La/Y, and high MgO) is precluded on three counts. First, magma mixing should produce straight arrays in binary plots. However, Al<sub>2</sub>O<sub>3</sub>, the light rare earth elements and Sr display inflections at around 60 wt.% SiO<sub>2</sub> (Fig. 2b,d,f). Second, adakitic rocks are more common in the west of the peninsula while arc lavas occur in the east (Fig 3e and f). This observation conflicts with models based on theory or experimental data, which predict that partial melting of subducted crust should occur closer to the trench than the fluid-fluxed melting of the mantle wedge which produces typical arc magmas [8-10,36]. Finally, if the adakitic melts had interacted with arc magma then they should possess isotopic characteristics intermediate between those of the slab and arc lavas. Figure 5 demonstrates that the adakites are entirely distinct from Philippine Sea Plate crust and that their <sup>87</sup>Sr/<sup>86</sup>Sr and <sup>143</sup>Nd/<sup>144</sup>Nd ratios are very similar to Pliocene arc rocks. In making this comparison we have deliberately chosen data for basaltic rocks from the shallow crust of the Philippine Sea Plate that were not acid-leached prior to analysis, as these will most accurately represent the subducted crust. Using analyses of acid-leached basalt from the slab would only increase the discrepancy between the slab and adakitic compositions. Furthermore, <sup>87</sup>Sr/<sup>86</sup>Sr tends to decrease with depth in altered oceanic crust [37,38]. If slab melts are produced as average of melt fractions from the upper and lower crust [39] then they should have even lower <sup>87</sup>Sr/<sup>86</sup>Sr than the shallowest lavas. The isotopic differences between adakites and slab suggest that the former are not derived



directly from the latter. Instead, both Pliocene and Pleistocene rocks are displaced to high  $^{87}\text{Sr}/^{86}\text{Sr}$  relative to the  $^{143}\text{Nd}/^{144}\text{Nd}$  of the upper mantle (Fig. 5). In conjunction with the similarities of all trace element ratios, except those involving Y and the HREEs (Fig. 4a), this suggests both the adakitic and arc suites are ultimately derived from similar sources.

The second means of changing slab melt composition is assimilation of rocks from the mantle or crust. With respect to  $\text{SiO}_2$  the Surigao adakites possess high mg-numbers (Table 1) and are relatively rich in elements that are abundant in peridotite, such as MgO (Fig. 2a) and Ni. Similar characteristics have been interpreted as evidence of melt-mantle interaction in other adakitic suites [14,40]. Laboratory experiments demonstrate that variable interaction between slab-derived adakite and peridotite should produce suites of silicic magma in which  $\text{SiO}_2$  correlates positively with  $\text{Al}_2\text{O}_3$  and negatively with  $\text{Na}_2\text{O}$  or  $\text{K}_2\text{O}$ . This is because assimilation involves precipitation of orthopyroxene, depleting the melt in silica but enriching it in incompatible elements [41]. Neither of these relationships is observed in the Surigao adakites. Indeed there is a strong positive correlation between silica and  $\text{K}_2\text{O}$  (Fig. 2c). Furthermore, reaction between silicic melt and peridotite can modify trace element concentrations in the hybridized melts but has a negligible effect on ratios of incompatible trace elements [41]. Therefore, even if such a reaction had modified  $\text{SiO}_2$  and MgO in some Surigao adakites it would not have a significant effect on key ratios such as  $\text{Sr}/\text{Y}$  and  $\text{La}/\text{Y}$ . The correlations in Figure 3a-c suggest that the major and trace element systematics of the adakites result from a common process, and not one that can leave little imprint on incompatible trace element ratios.

(Figure 5 close to this section)

Sr and Nd isotopic data are also inconsistent with a major role for interaction between slab melt and mantle wedge. Mantle peridotite beneath the Philippines has lower  $^{87}\text{Sr}/^{86}\text{Sr}$  than the subducted Philippine Sea Plate at similar  $^{143}\text{Nd}/^{144}\text{Nd}$ . Slab melts interacting with

this wedge would have their compositions driven towards lower  $^{87}\text{Sr}/^{86}\text{Sr}$ , i.e. away from those of Surigao adakites (Fig. 5). The mantle wedge may have been modified by slab-derived fluids during earlier phases of subduction, but it is unlikely that any peridotitic lithology would contain sufficient Sr to buffer  $^{87}\text{Sr}/^{86}\text{Sr}$  in the adakites, which are particularly rich in Sr (Fig. 2d). Any contaminant would require even higher Sr contents (>1200ppm) to influence  $^{87}\text{Sr}/^{86}\text{Sr}$  whilst having a negligible effect on other aspects of melt chemistry (Fig. 4a).

There is insufficient control on the isotopic composition of the Surigao basement to unequivocally dismiss the possibility that these isotope ratios have been modified by assimilation of crustal rocks. However,  $^{87}\text{Sr}/^{86}\text{Sr}$  and  $^{143}\text{Nd}/^{144}\text{Nd}$  do not correlate with  $\text{SiO}_2$  or  $\text{MgO}$  as would be predicted if crustal rocks were assimilated during differentiation. Furthermore, a contaminant with exceptionally high Sr, and Sr/Nd, would again be required to displace the isotope ratios of true slab melts away from those of the Philippine Sea Plate (Fig. 5).

#### *4.2 Variable Melting of the Philippine Sea Plate*

Low degree partial melts of hydrous basalt are rich in  $\text{SiO}_2$  and alkalis but poor in  $\text{MgO}$  and  $\text{FeO}$  [8,9]. As melting progresses silica and alkalis in the melt are diluted and the concentrations of ferromagnesian components increase. The Surigao suite displays these characteristics (Fig. 2a-c) so may result from variable degrees of partial melting of subducted basalt crust. However, the trace element and isotopic characteristics are not consistent with this origin. Basaltic rocks from the Philippine Sea Plate contain too little Sr to replicate the trace element variation observed in the Surigao suite. Even doubling the Sr content of the subducted basalt, to simulate seafloor alteration [8,38], results in Sr – Y variation unlike that observed at Surigao, or in any putative slab melts (short dashed line, Fig. 3d).

$^{87}\text{Sr}/^{86}\text{Sr}$  and  $^{143}\text{Nd}/^{144}\text{Nd}$  data also suggest that Surigao adakites were not generated by melting of Philippine Sea Plate crust. Variations in partial melting would have a negligible impact on the Sr and Nd isotopic ratios of the magmas produced. Instead of resembling the subducted slab, the Surigao adakites are most similar to Pliocene arc lavas (Section 4.1). Slab melting is also inconsistent with osmium isotope data [42]. The high Re/Os ratios of ocean floor basalt mean that, at more than 50 million years old, the Philippine Sea Plate crust should have developed extremely high  $^{187}\text{Os}/^{188}\text{Os}$ , which would also be passed on to slab melts. In fact, the majority of Surigao adakitic rocks possess  $^{187}\text{Os}/^{188}\text{Os}$  within the range of most island arc lavas [42].

#### *4.3 Adakite production from Arc Basalt*

Incompatible trace element ratios and isotopic characteristics of the Pleistocene rocks are similar to those of Pliocene arc lavas except for extreme depletion of Y and the HREEs (Fig. 4b). Y and the HREE are strongly correlated with  $\text{SiO}_2$  (Fig. 2e and 4b). Since, (i) magma mixing, assimilation of mantle or crust, and variable slab melting cannot satisfactorily explain the geochemistry of the adakites, and (ii) the isotopic data suggest that the Pliocene and Pleistocene rocks ultimately share a source in mantle wedge, we conclude that the adakitic signature of the Pliocene rocks was produced either by solid fractionation from arc magma, or by partial melting of arc magma that had completely solidified.

Plagioclase is by far the most abundant phenocryst in the adakitic rocks (25-50%), followed by hornblende (10-15%), with trace quantities of biotite, Fe-Ti oxide and clinopyroxene. Differentiation of an amphibole-dominated assemblage has been proposed as a mechanism to produce adakitic rocks on Camiguin Island, north of Mindanao [43], but a plagioclase-amphibole assemblage is unable to reproduce the trace element signature of the Surigao suite. In particular, removal of these phases would produce concave-upwards patterns between the middle and heavy rare earth elements (Fig. 4c) and

result in decreasing Dy/Yb with increasing SiO<sub>2</sub>. The increase of Dy/Yb with differentiation (Fig. 3c) requires that a phase with  $D_{Yb} > D_{Dy}$ , such as garnet, was involved in the development of the adakitic signature. The rare earth element patterns are consistent with fractionation of an assemblage containing clinopyroxene, orthopyroxene, garnet and amphibole in proportions similar to those crystallised in basaltic melt at 1.2GPa (Fig. 4c; [44]). Crystallisation of a small quantity of a light rare earth element-bearing phase, such as allanite, is also required to account for the depletion of La and Ce in the most silica-rich compositions (Figs. 2f and 4). Employing Pliocene arc lava as a starting composition suggests that 30% to 50% crystallisation of the high pressure assemblage is sufficient to produce the the Surigao suite (solid line, Fig. 3d).

Alternatively, adakitic rocks may be produced by remelting arc magma that solidified at depths where garnet was stable. Garnet and amphibole may crystallise from basaltic melt at high pressure (see above; [44]) or may develop during isobaric cooling of rocks emplaced slightly shallower than the depth where garnet becomes a liquidus phase [45]. As already noted, major element variations in the Surigao suite closely resemble andesite and dacite compositions generated in the laboratory by isobaric melting of hydrous metabasalt over a range of temperatures (Fig. 2a-c). Remelting arc lava, which contains more Sr (and other trace elements) than ocean floor basalt, produces a better fit to adakitic Sr - Y systematics than melting subducted ocean crust (long dash line, Fig. 3d) but requires extremely high degrees (>50%) of batch melting with residual garnet, amphibole and pyroxene (Fig. 3d). Such high degrees of melting may not generate adakite if garnet is a minor component of the source, c.f. [44], although there is experimental evidence that garnet is precipitated during low degrees of partial melting of hydrous basalt [8,9].

Castillo et al. [43] suggested that shallow, amphibole-dominated differentiation of arc lavas may generate some Philippine adakites. Remelting basaltic rock in thick, garnet-bearing crust has been proposed as a mechanism for generating adakitic magma from arc lithosphere [24-28]. Furthermore, garnet-, amphibole-bearing rocks have been

documented from basal sections of exhumed arc crust in the Aleutian arc and Kohistan, where Moho depths are estimated to have been  $\geq 30\text{km}$  [45-47]. The Surigao example is distinct from these models in revealing a strong garnet fractionation signature in magmatism associated with relatively thin arc crust. Therefore, our data require that adakitic compositions were generated as a result of basaltic melt crystallising within the mantle.

A case can be made, based on simple buoyancy arguments, that basaltic melt should not pond until it reaches the Moho, which in arcs with crust less than  $\sim 35\text{km}$  thick is too shallow for garnet crystallisation. However, Stratford and Stern [48] have imaged a strong seismic reflection and large drop in shear wave velocities at  $35\text{km}$  depth beneath the Taupo Volcanic Zone, New Zealand, where the crust is less than  $20\text{km}$  thick. They interpret this anomaly as a rising diapir of melt or a melt body trapped at a thermal boundary layer within the mantle. Basic arc magma provides the most likely candidate for a liquid body at this depth. A stalled diaper or melt body implies a mechanical and/or rheological barrier impeding further upward migration. Whether such a barrier lies within the arc lithosphere or marks a boundary between arc lithosphere and the underlying, convecting mantle is beyond the scope of this paper. However, the presence of substantial melt volumes within the shallow mantle wedge is consistent with the inferences from the Surigao geochemical data. At  $35\text{km}$  basaltic arc magma will crystallise a garnet-bearing assemblage. As discussed above this, in turn, will produce either silicic differentiated liquid with adakitic chemistry, or garnet-bearing mafic rock that could remelt to yield adakitic magma. In either case, during transport from the locus of crystallisation to the Moho the adakitic magma produced will have the opportunity to interact with mantle peridotite and acquire the elevated MgO, Ni and Cr concentrations and mg-numbers observed at Surigao and in other adakite suites [14].

#### 4.4 Adakite Production in the East Philippine Arc

Surigao's low-lying central plain (Fig. 1c) is a young rift or transtensional feature [32], so the temperature at any depth beneath the plain will be higher than at the same depth further west, on the rift margin (Fig. 6). The strong spatial control on the composition of Surigao adakites infers that the temperature of solid – melt equilibrium increases from the west coast to the central plain (Fig. 3e and f). Lower geothermal gradients at the rift margin may allow more extensive crystallisation of magma here than is the case in the central part of the rift. Similarly, if remelting is responsible for adakite generation, then melting temperatures will be higher beneath the central plain than at a similar depth beneath the rift margins (Fig. 6). Therefore, Pleistocene thinning of the overriding plate provides a single mechanism to produce both the rifted morphology of the peninsula and the geographic control on melt chemistry.

#### 4.5 Implications for Other Adakite Suites

Figure 6 summarises the mechanisms by which adakitic magma may be generated from arc basalt via crystallisation in the mantle wedge. This model has several important implications for petrogenesis of other adakitic magmatic suites.

(Figure 6 here)

First, adakitic magma can be derived from primitive arc magma, which is consistent with  $\delta^{18}\text{O}$  values of adakites. Oceanic lithosphere and sediment are very heterogeneous in  $\delta^{18}\text{O}$  [38,49-51], therefore, similar diversity would be predicted for melts derived from subducted slabs. However, adakitic rocks display a narrow range of  $\delta^{18}\text{O}$  values [39] extending only slightly higher than other subduction zone magmatic rocks [51-55]. Pyroxene, garnet and amphibole all have lower  $\delta^{18}\text{O}$  values than silicate melt with which they are in equilibrium [39,56]. Fractionation of these phases from magma, either by fractional crystallisation or partial melting, will produce a small increase in  $\delta^{18}\text{O}$  values of

the differentiated melts. This is more consistent with the limited oxygen isotopic variation in adakitic rocks than the fortuitous balance of sources required during melting of different slabs, each displaying its own diverse  $\delta^{18}\text{O}$  distribution.

If adakitic magma can be produced from any primitive arc melt, why are adakites not more common in more arcs? The architecture, rheology and thermal structure of a subduction zone will control the extent to which deep crystallisation may occur and be overprinted by later differentiation. In Surigao, deep differentiation is recorded while shallow crystallisation appears to have had a negligible impact on geochemistry of the adakites. In contrast, the Pliocene rocks are typical arc lavas and do not record deep processing. This difference could reflect changes in magma plumbing across the arc with deeper ponding favoured at greater distance from the arc. Alternatively, variations in the magma flux through the arc could be responsible. An extensive magma plumbing system in mature arc crust will decrease the probability that the signature of deep crystallisation will survive shallow crystallisation, magma mixing or interaction with the crust itself (Fig. 6). In eastern Mindanao (or, indeed much of the eastern Philippines) there is currently no active volcanism associated with subduction of the Philippine Sea Plate, suggesting low melt productivity from the mantle wedge. The Pleistocene adakites were the last magmatic event to affect Surigao and so represent a waning melt flux to the crust. Ephemeral magmatism during the earliest stages of magmatism may, therefore, be conducive to creating conditions under which adakitic magmas can avoid further differentiation, and loss of their distinctive character, at shallower levels. A component with “slab fusion” characteristics has also been invoked in rocks generated in the nascent Izu-Bonin arc [57] but may, in fact, represent remobilisation of basaltic rock emplaced during earlier phases of subduction [30,58]. If differentiation in the mantle is responsible for adakite generation, then the link between young subducted lithosphere and those adakites identified by Drummond and Defant [10] could reflect the thermal and dynamic effect of young slabs on the shallow mantle wedge and overriding lithosphere, rather than melting of the slab itself.

Second, our conclusions indicate that unusually hot subducted slabs are not a prerequisite of adakitic magmatism. The elegance of the original adakite model [10] was that each adakitic suite was linked by the common denominator of a young slab, which was inferred to be hot. Subsequent recognition of adakites associated with old slabs has broken this simple link but thrown up several exceptions to the rule. Each new case requires its own mechanism for heating old, cold slab to melting point. Two extreme examples of the absence of a hot slab are (i) genesis of adakitic magma where the slab is at very great depth [59], or (ii) genesis of adakites without any slab at the time of magmatism [60-62]. The alternative to having several exceptions to the rule is to have a new rule. Basalt that crystallises in the lithospheric mantle (Stage 2, Fig. 6) will remain part of the lithosphere until later perturbations of the geothermal gradient. Partial melting of arc basalt stored in the lithospheric mantle would still result in adakitic magma regardless of the geodynamic process causing the basalt to melt. Non-arc basalts have different bulk compositions, lower water contents and less distinctive trace element ratios than those produced in subduction zones. Therefore, their remobilisation from sites of deep crystallisation may not happen so readily or will not produce such distinctive partial melts as those with a subduction zone provenance.

Third, the shallow mantle may act as a staging-post for at least some primitive magmas in subduction zones [11,62,63]. Evidence for interaction between adakitic melt and peridotite has previously been used as evidence that adakites must originate beneath the Moho [13,14,22,40] and thus, by inference, in the subducted slab. However, Surigao demonstrates that adakitic magma produced from deeply crystallised basalt must still traverse the uppermost mantle wedge (Stage 3; Fig. 6). Our model infers that mafic to ultramafic cumulates can develop within the mantle. In turn, this provides a location where a primary basaltic magma flux into the base of arc lithosphere can generate a magmatic flux into the crust that has a more evolved bulk composition [11,44,46,62].



## 5. Summary

Pleistocene igneous rocks from Surigao record the development of adakitic melts from typical arc magma by fractionation of a garnet-bearing assemblage. Arc magma stalled within the mantle, either at the base of the arc lithosphere or at some rheological boundary in the shallow mantle wedge. Fractional crystallisation or remelting of the stalled material produced the adakites. Our data imply that (i) adakitic magmatism can occur in any subduction zone where “normal” arc magmatism occurs, (ii) adakitic magmatism does not require an unusually hot subducted slab, (iii) adakitic magmatism can be generated out-with active subduction zones, and (iv) that the mantle beneath subduction zones can play an important role in determining the nature of volcanic outputs and of crust produced by subduction zones. These conclusions have important implications for interpreting geodynamics of modern adakites and adakite-related rocks found in Archean terranes.

## Acknowledgements

The SE Asia Research Group, Royal Holloway University of London, funded much of this work. Dreher was supported by a Royal Society USA Research Fellowship. N. Ruelo and P. Waters of Anglo American (Minorco) helped arrange and conduct fieldwork in the Philippines. L. Whittaker prepared the rocks for major element analysis. T. Bostock and C. Ottley conducted ICP-MS analysis at Durham. Discussion with P.R. Castillo, J.P. Davidson, R. Hall, Y. Niu, D.G. Pearson and R.N. Thompson helped refine these ideas and anonymous comments during journal review helped improve the manuscript. We thank K. Farley for his editorial work.

## References

- [1] S.M. Peacock, T. Rushmer, A.B. Thompson, Partial melting of subducting oceanic crust, *Earth Planet. Sci. Lett.* 121 (1994) 227-244.

- 409 [2] M.W. Schmidt, S. Poli, Experimentally based water budgets for dehydrating slabs  
410 and consequences for arc magma generation, *Earth Planet. Sci. Lett.* 163 (1998)  
411 361-379.
- 412 [3] T.V. Gerya, D.A. Yuen, Rayleigh-Taylor instabilities from hydration and melting  
413 propel 'cold plumes' at subduction zones, *Earth Planet. Sci. Lett.* 212 (2003) 47-62.
- 414 [4] C.J. Hawkesworth, K. Gallagher, J.M. Hergt, F. McDermott, Mantle and slab  
415 contributions in arc lavas, *Ann. Rev. Earth Planet. Sci.* 21 (1993) 175-204.
- 416 [5] J.A. Pearce, D.W. Peate, Tectonic implications of the composition of volcanic arc  
417 magma, *Ann. Rev. Earth Planet. Sci.* 23 (1995) 251-285.
- 418 [6] J.P. Davidson, Decipheric mantle and crustal signatures in subduction zone  
419 magmatism, in: G.E. Bebout, D. Scholl, S. Kirby, J.P. Platt, (Eds.), *Subduction Top  
420 to Bottom*, AGU Monograph 96 (1996) 251-262.
- 421 [7] S. Turner, C.J. Hawkesworth, Constraints on the flux rates and mantle dynamics  
422 beneath island arc volcanoes from Tonga-Kermadec lava geochemistry. *Nature* 389  
423 (1997) 568-573.
- 424 [8] C. Sen, T. Dunn, Dehydration melting of a basaltic composition amphibolite at 1.5  
425 and 2.0 GPa: Implications for the origin of adakites. *Contrib. Mineral. Petrol.* 117  
426 (1994) 394-409.
- 427 [9] R.P. Rapp, E.B. Watson, Dehydration melting of metabasalt at 8-32 kbar:  
428 implications for continental growth and crust-mantle recycling, *J. Petrol.* 36 (1995)  
429 891-931.
- 430 [10] M.J. Defant, M.S. Drummond, Derivation of some modern arc magmas by melting  
431 of young subducted lithosphere, *Nature* 347 (1990) 662-665.
- 432 [11] P.D. Kelemen, 1995, Genesis of high Mg-Number andesites and the continental  
433 crust: *Contrib. Mineral. Petrol.* 120 (1995) 1-19.
- 434 [12] M.S. Drummond, M.J. Defant, P.K. Kepezhinskis, Petrogenesis of slab-derived  
435 trondhjemite-tonalite-dacite/adakite magmas, *Trans. R. Soc. Edinburgh* 87 (1996)  
436 205-215.
- 437 [13] H. Martin, Adakitic magmas: modern analogues of Archaean granitoids, *Lithos* 46  
438 (1999) 411-429.
- 439 [14] R.H. Smithies, The Archaean tonalite-trondhjemite-granodiorite (TTG) series is not  
440 an analogue of Cenozoic adakite, *Earth Planet. Sci. Lett.* 182 (2000) 115-125.
- 441 [15] D. Thieblemont, G. Stein, G.L. Lescuyer, Epithermal and porphyry deposits: the  
442 adakite connection, *Comptes Rendus de l'Academie des Sciences Serie II –  
443 Sciences de la Terre et des Planetes* 325 (1997) 103-109.
- 444 [16] F.G. Sajona, R. Maury, Association of adakites with gold and copper mineralization  
445 in the Philippines, *Comptes Rendus de l'Academie des Sciences Serie II – Sciences  
446 de la Terre et des Planetes*, 326 (1998) 27-34.
- 447 [17] R. Oyarzun, A. Márquez, J. Lillo, I. López, S. Rivera, Giant versus small porphyry  
448 copper deposits of Cenozoic age in northern Chile: adakitic versus normal calc-  
449 alkaline magmatism, *Mineralium Deposita* 36 (2001) 794-798.

- [18] J. Richards, Discussion on “Giant versus small porphyry copper deposits of Cenozoic age in northern Chile: adakitic versus normal calc-alkaline magmatism” by Oyarzun et al. (*Mineralium Deposita* 36: 794-798, 2001), *Mineralium Deposita* 37 (2002) 788-790.
- [19] O. Rabbia, L.B. Hernández, R.W. King, L. López-Escobar, Discussion on “Giant versus small porphyry copper deposits of Cenozoic age in northern Chile: adakitic versus normal calc-alkaline magmatism” by Oyarzun et al. (*Mineralium Deposita* 36: 794-798, 2001), *Mineralium Deposita* 37 (2002) 791-794.
- [20] R. Oyarzun, A. Marquez, J. Lillo, I Lopez, S. Rivera, Reply to discussion on “Giant versus small porphyry copper deposits of Cenozoic age in northern Chile: adakitic versus normal calc-alkaline magmatism” by Oyarzun R, Márquez A, Lillo J, López I, Rivera S (*Mineralium Deposita* 36: 794-798, 2001), *Mineralium Deposita* 37 (2002) 795-799.
- [21] F.G. Sajona, R.C. Maury, H. Bellon, J. Cotton, M.J. Defant, M. Pubellier, Initiation of subduction and the generation of slab melts in western and eastern Mindanao, Philippines, *Geology* 21 (1993) 1007-1110.
- [22] G.M. Yogodzinski, J.M. Lees, T.G. Churikova, F. Dorendorf, G. Wöerner, O.N. Volynets, Geochemical evidence for the melting of subducting oceanic lithosphere at plate edges, *Nature* 409 (2001) 500-504.
- [23] M.-A. Gutscher, R. Maury, J.-P. Eissen, E. Bourdon, Can slab melting be caused by flat subduction? *Geology* 28 (2000) 535-538.
- [24] M.P. Atherton, N. Petford, Generation of sodium-rich magmas from newly underplated basaltic crust, *Nature* 362 (1995) 144-146.
- [25] R.M. Conrey, P.R. Hooper, P.B. Larson, J. Chesley, J. Ruiz, Trace element and isotopic evidence for two types of crustal melting beneath a High Cascade volcanic center, Mt. Jefferson, Oregon: *Contrib. Mineral. Petrol.* 141 (2001) 710-732.
- [26] J.M. Garrison, J.P. Davidson, Dubious case for slab melting in the Northern volcanic zone of the Andes, *Geology* 31 (2003) 565-568.
- [27] Z.Q. Hou, Y.F. Gao, X.M. Qu, Z.Y. Rui, X.X. Mo, Origin of adakitic intrusives generated during mid-Miocene east-west extension in southern Tibet, *Earth Planet. Sci. Lett.* 220 (2004) 139-155.
- [28] O. Wang, F. McDermott, J.F. Xu, H. Bellon, Y.T. Zhu, Cenozoic K-rich adakitic volcanic rocks in the Hohxil area, northern Tibet: Lower-crustal melting in an intracontinental setting, *Geology* 33 (2005) 465-468.
- [29] M. Pubellier, R. Quebral, C. Rangin, B. Deffontaines, C. Muller, J. Butterlin, J. Manzano, The Mindanao collision zone: a soft collision event within a continuous Neogene strike-slip setting, *J. South East Asian Earth Sci.* 6 (1991) 239-248.
- [30] C.G. Macpherson, R. Hall, Tectonic setting of Eocene boninite magmatism in the Izu-Bonin-Mariana forearc, *Earth Planet. Sci. Lett.* 186 (2001) 215-230.
- [31] A.H.G. Mitchell, T.M. Leach, *Epithermal Gold in the Philippines: Island arc metallogeneses, geothermal systems and geology*, Academic Press, London, (1991) pp. 457.

- [32] R.D. Quebral, M. Pubellier, C. Rangin, The onset of movement on the Philippine Fault in eastern Mindanao: a transition from a collision to a strike-slip environment, *Tectonics* 15 (1996) 713-726.
- [33] C.G. Macpherson, R. Hall, Timing and tectonic controls in the evolving orogen of SE Asia and the western Pacific and some implications for ore generation, in: D.J. Blundell, F. Neubauer, F. von Quandt (Eds.), *The Timing and Location of Major Ore Deposits in an Evolving Orogen*, Geol. Soc. Long. Spec. Publ. 204 (2002) 49-67.
- [34] M. Pubellier, C. Monnier, R. Maury, R. Tamayo, Plate kinematics, origin and tectonic emplacement of supra-subduction ophiolites in SE Asia, *Tectonophysics* 392 (2004) 9-36.
- [35] S.B. Mukasa, R. McCabe, J.B. Gill, Pb-isotopic composition of volcanic rocks in the west and east Philippine island arcs: presence of the DUPAL isotopic anomaly, *Earth Planet. Sci. Lett.* 84 (1987) 153-164.
- [36] M.J. Defant, M.S. Drummond, Mount St. Helens: Potential example of the partial melting of the subducted lithosphere in a volcanic arc, *Geology* 21 (1993) 547-550.
- [37] M.T. McCulloch, R.T. Gregory, G.J. Wasserburg, H.P. Taylor, A neodymium, strontium, and oxygen isotopic study of the Cretaceous Samail Ophiolite and implications for the petrogenesis and seawater-hydrothermal alteration of oceanic crust, *Earth Planet. Sci. Lett.* 46 (1980) 201-211.
- [38] H. Staudigel, G.R. Davies, S.R. Hart, K.M. Marchant, B.M. Smith, Large scale isotopic Sr, Nd and oxygen isotopic anatomy of altered oceanic crust: DSDP/ODP sites 417/418, *Earth Planet. Sci. Lett.* 130 (1995) 169-185.
- [39] I.N. Bindeman, J.M. Eiler, G.M. Yogodzinski, Y. Tatsumi, C.R. Stern, T.L. Grove, M. Portnyagin, K. Hoernle, L.V. Danyushevsky, Oxygen isotope evidence for slab melting in modern and ancient subduction zones, *Earth Planet. Sci. Lett.* 235 (2005) 480-496.
- [40] R.W. Kay, Aleutian magnesian andesites: melts from subducted Pacific Ocean crust, *J. Volcanol. Geotherm. Res.* 4 (1978) 117-132.
- [41] R.P. Rapp, N. Shimizu, M.D. Norman, Reaction between slab-derived melts and peridotite in the mantle wedge: experimental constraints at 3.8 GPa, *Chem. Geol.* 160 (1999) 335-356.
- [42] S.T. Dreher, C.G. Macpherson, D.G. Pearson, J.P. Davidson, Re-Os isotope studies of Mindanao adakites: implications for sources of metals and melts, *Geology* 33 (2005) 957-960.
- [43] P.R. Castillo, P.E. Janney, R.U. Solidum, Petrology and geochemistry of Camiguin Island, southern Philippines: insights to the source of adakites and other lavas in a complex arc setting, *Contrib. Mineral. Petrol.* 134 (1999) 33-51.
- [44] O. Müntener, P.B. Kelemen, T.L. Grove, The role of H<sub>2</sub>O during crystallization of primitive arc magmas under uppermost mantle conditions and genesis of igneous pyroxenites: an experimental study, *Contrib. Mineral. Petrol.* 141 (2001) 643-658.

- [45] S.M. DeBari, R.G. Coleman, Examination of the deep levels of an island arc: evidence from the Tonisa ultramafic-mafic assemblage, Tonisa, Alaska, *J. Geophys. Res.* 94, (1989) 4373-4391.
- [46] S.M. DeBari, N.H. Sleep, High-Mg, low-Al bulk composition of the Talkeetna island arc, Alaska: implications for primary magmas and the nature of arc crust, *Geol. Soc. Am. Bull.* 103 (1991) 37-47.
- [47] P.J. Treloar, M.G. Pettersen, M. Quasim Jan, M.A. Sullivan, A re-evaluation of the stratigraphy and evolution of the Kohistan arc sequence, Pakistan Himalaya: implications for magmatic and tectonic arc-building processes. *J. Geol. Soc. Lond.* 153 (1996) 681-693.
- [48] W.R. Stratford, T.A. Stern, Strong seismic reflections and melts in the mantle of a continental back-arc basin, *Geophys. Res. Lett.* 31 (2004) art. no. L06622.
- [49] J.C. Alt, K. Muehlenbachs, J. Honnorez, An oxygen isotope profile through the upper kilometer of the oceanic crust, DSDP Hole 504B, *Earth Planet. Sci. Lett.* 80 (1986) 217-227.
- [50] K. Muehlenbachs, Alteration of the oceanic crust and the  $^{18}\text{O}$  history of seawater, in: J.W. Valley, H.P. Taylor, J.R. O'Neil (Eds), *Stable isotopes in high temperature geological processes*, *Min. Soc. Am. Rev.* 16 (1986) 425-444.
- [51] C.G. Macpherson, J.A. Gamble, D.P. Matthey, Oxygen isotope geochemistry of lavas from an oceanic to continental arc transition, Kermadec – Hikurangi margin, SW Pacific, *Earth Planet. Sci. Lett.* 160 (1998) 609-621.
- [52] M.F. Thirlwall, A.M. Graham, R.J. Arculus, R.S. Harmon, C.G. Macpherson, Resolution of the effects of crustal assimilation, sediment subduction, and fluid transport in island arc magmas: Pb-Sr-Nd-O isotope geochemistry of Granada, Lesser Antilles. *Geochim. Cosmochim. Acta* 60 (1996) 4785-4810.
- [53] C.G. Macpherson, D.P. Matthey, Oxygen isotope variations in Lau Basin lavas, *Chem. Geol.* 144 (1998) 177-194.
- [54] C.G. Macpherson, D.R. Hilton, D.P. Matthey, J.M. Sinton, Evidence for an  $^{18}\text{O}$ -depleted mantle plume from contrasting  $^{18}\text{O}/^{16}\text{O}$  ratios of back-arc lavas from the Manus Basin and Mariana Trough, *Earth Planet. Sci. Lett.* 176 (2000) 171-183.
- [55] J.M. Eiler, A. Crawford, T. Elliott, K.A. Farley, J.W. Valley, E.M. Stolper, Oxygen isotope geochemistry of oceanic arc lavas, *J. Petrol.* 41 (2000) 229-256.
- [56] J.-F. Zheng, Calculation of oxygen isotope fractionation in metal oxides, *Geochim. Cosmochim. Acta* 55 (1991) 2299-2307.
- [57] J.A. Pearce, M.F. Thirlwall, G. Ingram, B.J. Murton, R.J. Arculus, S.R. van der Laan, Isotopic evidence for the origin of the boninites and related rocks drilled in the Izu-Bonin (Ogasawara) forearc, leg 125, in: P. Fryer, J.A. Pearce, L. Stokking, et al., (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results 125*, Ocean Drilling Program, College Station (1992) 623-659.
- [58] R. Hall, Cenozoic geological and plate tectonic evolution of SE Asia and the SW Pacific: computer-based reconstructions, model and animations, *J. Asian Earth Sci.* 20 (2002) 353-431.

- 575 [59] F.G. Sajona, R.C. Maury, M. Pubellier, J. Leterrier, H. Bellon, J. Cotton, Magmatic  
576 source enrichment by slab-derived melts in a young post-collisional setting, central  
577 Mindanao (Philippines), *Lithos*, 54 (2000) 173-206.
- 578 [60] R.W. Johnson, D.E. Mackenzie, I.E.M. Smith, Delayed partial melting of  
579 subduction-modified mantle in Papua New Guinea, *Tectonophys.* 46 (1978) 197-  
580 216.
- 581 [61] I.E.M. Smith, S.R. Taylor, R.W. Johnson, REE-fractionated trachytes and dacites  
582 from Papua New Guinea and their relationship to andesite petrogenesis, *Contrib.*  
583 *Mineral. Petrol.* 69 (1979) 227-233.
- 584 [62] G. Rogers, A.D. Saunders, D.J. Terrell, S.P. Verma, G.F. Marriner, Geochemistry  
585 of Holocene volcanic rocks associated with ridge subduction in Baja California,  
586 Mexico, *Nature* 315 (1985) 389-392.
- 587 [63] P.B. Kelemen, G.M. Yogodzinski, D.W. Scholl, Along-strike variation in lavas of  
588 the Aleutian arc: implications for the genesis of high Mg# andesite and continental  
589 crust, in: J. Eiler, (Ed.), *Inside the Subduction Factory*, Geophysical Monograph  
590 138, American Geophysical Union, Washington D.C. (2003) 223-246.
- 591 [64] P.B. Kelemen, K.T.M. Johnson, R.J. Kinzler, A.J. Irving, High-field-strength  
592 element depletions in arc basalts due to mantle-magma interaction, *Nature* 345  
593 (1996) 521-524.
- 594 [65] R. Hickey-Vargas, Origin of the Indian Ocean-type isotopic signature in basalts from  
595 Philippine Sea Plate spreading centres: An assessment of local versus large-scale  
596 processes, *J. Geophys. Res.* 103 (1998) 20963-20979.
- 597 [66] S.-S. Sun, W.F. McDonough, Chemical and isotopic systematics of oceanic basalts:  
598 implications for mantle compositions and processes, in: A.D. Saunders, M.J. Norry  
599 (Eds.) *Magmatism in the Ocean Basins*. *Geol. Soc. Spec. Publ.* 42 (1989) 313-345.
- 600 [67] H. Martin, Petrogenesis of Archean trondhjemites, tonalites, and granodiorites from  
601 eastern Finland: major and trace element geochemistry, *J. Petrol.* 28 (1987) 921-  
602 953.
- 603 [68] R. Hickey-Vargas, Origin of the Indian Ocean-type isotopic signature in basalts  
604 from Philippine Sea plate spreading centres: An assessment of local versus large-  
605 scale processes, *J. Geophys. Res.* 103 (1998) 20963-20979.
- 606 [69] C.G. Macpherson, S. Dreher, M.F. Thirlwall, Adakites without slab melting,  
607 Mindanao, the Philippines, *Geophys. Res. Abstracts*, 7 (2005) 07201.
- 608 [70] L. Bodri, B. Bodri, Numerical investigation of tectonic flow in island-arc areas.  
609 *Tectonophys.* 50 (1978) 163-175.
- 610 [71] A. Rowland, J.H. Davies, Buoyancy rather than rheology controls the thickness of  
611 the overriding mechanical lithosphere at subduction zones, *Geophys. Res. Lett.* 26  
612 (1999) 3037-3040.
- 613 [72] Y. Fukahata, M. Matsu'ura, Effects of active crustal movements on thermal  
614 structure in subduction zones, *Geophys. J. Int.* 141 (2000) 271-281.

## Figure Captions

Figure 1. (a) Philippine archipelago showing location of the Surigao peninsula on Mindanao. Plate boundaries after Hall [58]. PF is the Philippine Fault. (b) Sampling locations of Pleistocene, late Miocene to Pliocene and Paleogene volcanic rocks. Dashed line is the Philippine Fault (PF) and solid, barbed lines are Miocene reverse faults that have been reactivated in the opposite sense. The low lying region between these faults and the PF is an extensional or transtensional basin [32]. (c) Radar image of Surigao topography viewed from SSW, with the coast and shore of Lake Mainit outlined in white.

Figure 2. Variation of selected major and trace element concentrations of Surigao magmatic rocks. Plots of (a) MgO, (b)  $\text{Al}_2\text{O}_3$ , (c)  $\text{K}_2\text{O}$ , (d) Sr, (e) Y and (f) La versus  $\text{SiO}_2$ . In (a)-(c) compositions of high pressure, synthetic partial melts of hydrous metabasalt [9] are shown for comparison.

Figure 3. (a) Sr/Y, (b) La/Y (normalised to N-MORB), and (c) Dy/Yb versus  $\text{SiO}_2$  in Surigao magmatic rocks. (d) Sr/Y versus Y showing fields of adakites and island arc andesites, dacites and rhyolites [10]. The solid line illustrates fractional crystallisation of a high pressure mineral assemblage (see caption to Fig. 4) from basaltic melt initially containing 555ppm Sr and 26.5ppm Y (357460). Tick marks indicate the fraction of melt remaining. Long dashed line shows partial melting of a rock with the same initial Sr and Y as 357460. Short dashed line is partial melting of basalt from DSDP site 291 on the Philippine Sea Plate [65] with 24.7ppm Y and its Sr content doubled to 218ppm to simulate seafloor alteration. Tick marks indicate the extent of partial melting, which were calculated using partition coefficients from [8] and a residual mineralogy of 50% garnet, 50% clinopyroxene [9]. (e)  $(\text{La/Y})_n$ , and (f)  $\text{SiO}_2$  versus longitude, which is a measure of depth to the subducted slab beneath Surigao.

Figure 4. (a) Incompatible trace element concentrations of Pliocene and Pleistocene andesites from Surigao with similar  $\text{SiO}_2$  contents (60.5 wt.%), normalised to N-MORB

[66]. (b) Chondrite-normalised rare earth element concentrations for Pleistocene adakitic rocks from Surigao showing progressive depletion in the heavy rare earth elements with increasing SiO<sub>2</sub>. (c) Chondrite-normalised rare earth element concentrations of liquid evolved from the lowest-SiO<sub>2</sub> (56.7 wt.%) adakite by fractional crystallisation at low and high pressures. Low pressure assemblage is plagioclase, amphibole and FeTi-oxide in the proportions 74.3 : 21.5 : 4.2 based on phenocrysts in the Pleistocene rocks. High pressure assemblage is clinopyroxene, orthopyroxene, garnet, amphibole and allanite in the proportions 52.8 : 17.2 : 12.3 : 17.4 : 0.2, based on the equilibrium assemblage in experimental basaltic melt containing 56.5 wt.% SiO<sub>2</sub> at 1.2GPa [44], but with the garnet fraction reduced by 20%. Partition coefficients from [8,67].

Figure 5. <sup>143</sup>Nd/<sup>144</sup>Nd versus <sup>87</sup>Sr/<sup>86</sup>Sr of Pliocene and Paleogene magmatic rocks from Surigao. Altered basalt from the subducting Philippine Sea Plate [68] and the Indian Ocean MORB [35] to represent the mantle wedge are shown for comparison. Surigao data from this work and [68].

Figure 6. Schematic illustration of adakite production by deep differentiation in an arc with thin crust. The slab is subducted beneath overriding arc lithosphere and induces corner flow (convection) in the mantle wedge. Dashed lines are schematic isotherms (for relatively low (L), medium (M) and high (H) temperature) illustrating that at any depth the shallow mantle is hottest where the arc lithosphere is thinnest [after 2,68-70]. Numbers in the vertical column refer to the flow diagram on the right, which summarises the possible mechanisms identified for generating adakitic melt without slab melting. Stage 1; genesis of primitive arc basalt. Stage 2; high pressure processing of basalt yields adakitic magma either directly, by fractional crystallisation, or indirectly, by remelting crystallised basaltic rock. Stage 3; interaction between adakitic magma and mantle peridotite. Stage 4; low-pressure crystallisation. Where the crust is thick Stage 2 can occur above the Moho and Stage 3 would be bypassed [24-26]. In mature arcs with a high magma flux to the crust Stage 4 will obscure or obliterate adakitic chemistry.






Figure 1




Figure 2




Figure 3

672

673





Figure 4

674

675



676

677




Figure 6